For Research Use Only

Pinin Recombinant antibody

Catalog Number:86292-1-RR



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number:

86292-1-RR BC062602 Protein A purification

GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000 $\mu g/ml$ by 5411 251028A12

Nanodrop: **UNIPROT ID:** Recommended Dilutions: Source Q9H307 WB: 1:5000-1:50000 Rabbit IF/ICC: 1:500-1:2000 Full Name:

Isotype: pinin, desmosome associated protein

IgG Calculated MW:

Immunogen Catalog Number: 82 kDa AG38940 Observed MW:

Applications

Tested Applications: Positive Controls:

WB, IF/ICC, ELISA WB: HeLa cells, HepG2 cells, HEK-293 cells, LO2 cells,

Species Specificity: DU 145 cells

human IF/ICC: HepG2 cells,

140 kDa

Background Information

PNN, also named as DRS or 140 kDa nuclear and cell adhesion-related phosphoprotein, is a 717 amino acid protein, which belongs to the pinin family. PNN localizes in the plasma membrane and is expressed in placenta, lung, liver, kidney, pancreas, spleen, thymus, prostate, testis, ovary, small intestine, colon, heart, epidermis, esophagus, brain and smooth and skeletal muscle. PNN is expressed strongly in melanoma metastasis lesions and advanced primary tumors. PNN as a transcriptional activator binds to the E-box 1 core sequence of the E-cadherin promoter gene. PNN is involved in the establishment and maintenance of epithelia cell-cell adhesion. PNN is a potential tumor suppressor for renal cell carcinoma. The calculated molecular weight of PNN is 82 kDa (native form), but the postmodified protein is 140 kDa (phospho form of isoform 1).

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 86292-1-RR (PNN antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using PNN antibody (86292-1-RR, Clone: 251028A12) at dilution of 1:1000 and Coralite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).