For Research Use Only

WBSCR17 Recombinant antibody

Catalog Number:86351-1-RR



Basic Information

Catalog Number: GenBank Accession Number:

86351-1-RR BC069624 GeneID (NCBI): Size: 100ul , Concentration: 1000 $\mu g/ml$ by 64409

Nanodrop: **UNIPROT ID:** Q6IS24

Rabbit Full Name:

Isotype: Williams-Beuren syndrome IgG chromosome region 17

Immunogen Catalog Number: Calculated MW: AG15792 598 aa. 68 kDa

Observed MW: 70~90 kDa

Applications

Tested Applications:

WB, ELISA

Species Specificity: human, mouse, rat

Positive Controls:

WB: mouse brain tissue, rat brain tissue, fetal human

Purification Method:

Protein A purification

Recommended Dilutions:

WB: 1:1000-1:6000

CloneNo.:

250868B10

brain tissue

Background Information

WBSCR17, also known as GALNT17, which encodes a brain-expressed N-acetylgalactosaminyl transferase (GalNAcT), is located at the distal edge of a region that is commonly deleted or duplicated in Williams Beuren Syndrome (WBS), a developmental disorder with motor and coordination problems, impaired visuospatial memory, and abnormal social interaction (PMID: 31554716). WBSCR17 loss-of-function has significant effects on cerebellar development, and is associated with phenotypes including developmental delay, deficits in motor coordination, reduced exploratory activity, and impaired social behavior (PMID: 22787146). With the calculated molecular mass of recombinant WBSCR17 being 68 kDa, the 70-90-kDa glycoproteins could also be detected due to posttranslational modifications (PMID: 22787146).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 86351-1-RR (WBSCR17 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLL) kinetic assays of 86351-1-RR against Human WBSCR17 were performed. The affinity constant is 1.25 nM.