

For Research Use Only

BMP9 Recombinant monoclonal antibody

Catalog Number: 86411-1-RR



Basic Information

Catalog Number: 86411-1-RR	GenBank Accession Number: NM_019506.4	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 12165	CloneNo.: 250512B9
Source: Rabbit	UNIPROT ID: Q9WV56	Recommended Dilutions: WB: 1:1000-1:4000
Isotype: IgG	Full Name: growth differentiation factor 2	
Immunogen Catalog Number: EG2482	Calculated MW: 48 kDa	
	Observed MW: 33 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : mouse liver tissue, rat liver tissue, pig liver tissue, mouse brain tissue, HepG2 cells, L02 cells
Species Specificity: human, mouse, rat, pig	

Background Information

Bone morphogenetic protein-9 (BMP9, also termed growth-differentiation factor-2, GDF-2) is a secreted signaling molecule that belongs to the transforming growth factor- β (TGF- β) super-family. First identified in fetal mouse liver cDNA libraries, BMP9 is predominantly produced by hepatic stellate cells and circulates at relatively high concentrations in adult plasma. BMP9 is synthesized as a precursor protein (pre-proBMP9) composed of 429 amino acids (aa) that includes a 22 aa signal peptide, a 297 aa prodomain (33 kDa) and a 110 aa mature protein (12.5 kDa). It is then cleaved by serine endoproteases, leading to a short dimeric mature form of 25 kDa, and the prodomain which can remain noncovalently associated with the mature short form, giving a size of about 100 kDa (PMID:21710321).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 86411-1-RR (Gdf2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.