

For Research Use Only

BAF170 Recombinant antibody

Catalog Number: 86473-3-RR



Basic Information

Catalog Number: 86473-3-RR	GenBank Accession Number: BC013045	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 6601	CloneNo.: 251330D11
Source: Rabbit	UNIPROT ID: Q8TAQ2	Recommended Dilutions: WB: 1:5000-1:50000
Isotype: IgG	Full Name: SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily c, member 2	
Immunogen Catalog Number: AG2634	Calculated MW: 1214 aa, 132 kDa	
	Observed MW: 170 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : LNCaP cells, A431 cells, HeLa cells, RAW 264.7 cells, C6 cells, NIH/3T3 cells, PC-12 cells
Species Specificity: human, mouse, rat	

Background Information

BAF170, also named as SWI/SNF complex 170 kDa subunit or SMARCC2, is a 1214 amino acid protein, which belongs to the SMARCC family. BAF170 is ubiquitously expressed. BAF170 is involved in transcriptional activation and repression of select genes by chromatin remodeling. BAF170 may be required for CoREST dependent repression of neuronal specific gene promoters in non-neuronal cells. BAF170 belongs to the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin remodeling complex (nBAF complex). The calculated molecular weight of BAF170, is 133 kDa, but modified BAF170 is about 170 kDa. (PMID: 8804307)

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 86473-3-RR (BAF170 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 86473-3-RR against Human BAF170 were performed. The affinity constant is 4.64 nM.