For Research Use Only

IL-10 Recombinant antibody, PBS Only (Capture)

86547-2-PBS

Nanodrop:

Catalog Number:86547-2-PBS



Basic Information

Catalog Number:

GenBank Accession Number:

NM_000572.3

GeneID (NCBI):

100ug, Concentration: 1 mg/ml by

UNIPROT ID: P22301

Rabbit Full Name: Isotype: interleukin 10 IgG Calculated MW:

Immunogen Catalog Number: 21 kDa

EG3296

Purification Method: Protein A purification

CloneNo.: 250059F11

Applications

Tested Applications:

Sandwich ELISA, Indirect ELISA, Sample test

Species Specificity:

Product Information

86547-2-PBS targets IL-10 as part of a matched antibody pair:

MP02490-1: 86547-2-PBS capture and 86547-1-PBS detection (validated in Sandwich ELISA)


Unconjugated rabbit recombinant monoclonal antibody in PBS only (BSA and azide free) storage buffer at a concentration of 1 mg/mL, ready for conjugation. Created using Proteintech's proprietary in-house recombinant technology. Recombinant production enables unrivalled batch-to-batch consistency, easy scale-up, and future security of supply.

This conjugation ready format makes antibodies ideal for use in many applications including: ELISAs, multiplex assays requiring matched pairs, mass cytometry, and multiplex imaging applications. Antibody use should be optimized by the end user for each application and assay.


Storage

Storage: Store at -80°C. Storage Buffer: PBS only, pH7.3


Selected Validation Data



Sandwich ELISA standard curve of MP02490-1, Human IL-10 Recombinant Matched Antibody Pair-PBS only. 86547-2-PBS was coated to a plate as the capture antibody and incubated with serial dilutions of standard Eg3296. 86547-1-PBS was HRP conjugated as the detection antibody. Range: 31.3-2000 pg/mL



Human peripheral blood mononuclear cells (PBMC) were cultured unstimulated or stimulated with 10 µg/mL PHA for 3 days. The mean IL-10 concentration was determined to be 169.2 pg/mL in unstimulated PBMC supernatant, 11,597.9 pg/mL in PHA stimulated PBMC supernatant.



Plasma of fifteen individual healthy human donors was measured. The IL-10 concentration of detected samples was determined to be 48.7 pg/mL with a range of 28.0-94.9 pg/mL