

For Research Use Only

CST4 Recombinant monoclonal antibody

Catalog Number:86955-1-RR



Basic Information

Catalog Number: 86955-1-RR	GenBank Accession Number: NM_001899.2	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 1472	CloneNo.: 252073B11
Source: Rabbit	UNIPROT ID: P01036	Recommended Dilutions: WB: 1:5000-1:50000
Isotype: IgG	Full Name: cystatin S	
Immunogen Catalog Number: EG3098	Calculated MW: 16 kDa	
	Observed MW: 16 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : human saliva,
Species Specificity: human	

Background Information

Cystatins are a family of inhibitors of cysteine peptidases that comprises the salivary cystatins D and S-type (including CST5 and CST4,1,2) and cystatin C-type (CST3) (PMID: 25329717). The D and S-type cystatin have also been detected in seminal plasma, tears, and tracheobronchial fluid but not in other fluids and secretions where cystatin C is found. And the CST4 was expressed in the serous acinar and demilune cells of the human submandibular gland and at lower levels in the serous acini of the parotid gland (PMID:11879580). CST4 specifically combines with cysteine protease to regulate its activity, thus preventing hydrolysis of the extracellular matrix. And some studies propose that CST4 might be a biomarker for gastrointestinal cancer (PMID:29218251; PMID:29636621).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human saliva were subjected to SDS PAGE followed by western blot with 86955-1-RR (CST4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 86955-1-RR against Human CST4 were performed. The affinity constant is 0.257 nM.