

For Research Use Only

NRAC Recombinant monoclonal antibody

Catalog Number: 87528-1-RR



Basic Information

Catalog Number: 87528-1-RR	GenBank Accession Number: NM_001008404.2	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 400258	CloneNo.: 252953C8
Source: Rabbit	UNIPROT ID: Q8N912	Recommended Dilutions: WB: 1:1000-1:4000
Isotype: IgG	Full Name: chromosome 14 open reading frame 180	
Immunogen Catalog Number: EG6025	Calculated MW: 18 kDa	
	Observed MW: 18 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : human heart tissue,
Species Specificity: human	

Background Information

NRAC (Nutritionally Regulated Adipose and Cardiac enriched protein, with the human homologous gene designated as C14ORF180) is a recently identified adipocyte-specific double-pass transmembrane protein. Both its N- and C-termini are located intracellularly, containing only a short extracellular loop and no known enzymatic domains. Its expression is significantly induced during the differentiation of white and brown adipocytes and is regulated by nutritional status: fasting and obesity both downregulate its mRNA levels, suggesting its role as a "fat sensor" in energy homeostasis. Mechanistically, NRAC forms a trimeric complex via its first transmembrane domain with the fatty acid transporter core protein CD36 and the membrane microdomain protein caveolin-1. When extracellular fatty acid concentrations increase, NRAC undergoes ubiquitination and internalization, causing CD36 to dissociate from caveolin-1 and switch to clathrin-mediated endocytosis. This process enhances fatty acid uptake, leading to adipocyte hypertrophy and increased overall adipose mass.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human heart tissue were subjected to SDS PAGE followed by western blot with 87528-1-RR (NRAC antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 87528-1-RR against Human NRAC were performed. The affinity constant is 0.801 nM.