

For Research Use Only

FST Recombinant monoclonal antibody, PBS Only (Capture)

Catalog Number: 87548-3-PBS



Basic Information

Catalog Number: 87548-3-PBS	GenBank Accession Number: NM_013409.3	Purification Method: Protein A purification
Size: 100ug, Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 10468	CloneNo.: 252591H1
Source: Rabbit	UNIPROT ID: P19883-1	
Isotype: IgG	Full Name: follistatin	
Immunogen Catalog Number: EG2981	Calculated MW: 38 kDa	

Applications

Tested Applications:
Sandwich ELISA, Indirect ELISA

Species Specificity:
human

Product Information

87548-3-PBS targets FST as part of a matched antibody pair:

MP03168-1: 87548-3-PBS capture and 87548-2-PBS detection (validated in Sandwich ELISA)

Unconjugated rabbit recombinant monoclonal antibody in PBS only (BSA and azide free) storage buffer at a concentration of 1 mg/mL, ready for conjugation. Created using Proteintech's proprietary in-house recombinant technology. Recombinant production enables unrivalled batch-to-batch consistency, easy scale-up, and future security of supply.

This conjugation ready format makes antibodies ideal for use in many applications including: ELISAs, multiplex assays requiring matched pairs, mass cytometry, and multiplex imaging applications. Antibody use should be optimized by the end user for each application and assay.

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS only, pH7.3


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HepG2 (human hepatocellular carcinoma cells) were cultured in DMEM supplemented with 10% fetal bovine serum, 2.5 mM L-glutamine, 100 U/mL penicillin, and 100 µg/mL streptomycin sulfate. An aliquot of the cell culture supernate was removed, assayed for human Follistatin, and measured 3.83 ng/mL.



PC-3 cells were cultured unstimulated or stimulated with 10 ng/mL TNF-A for 2 days. The mean Follistatin concentration was determined to be 5.66 ng/mL in unstimulated cell culture supernatant, 7.48 ng/mL in stimulated cell culture supernatant.



Plasma of eight individual healthy human donors was measured. The human Follistatin concentration of detected samples was determined to be 1.69 ng/mL with a range of 1.14 - 3.10 ng/mL.



Sandwich ELISA standard curve of MP03168-1, Human Follistatin Recombinant Matched Antibody Pair - PBS only. 87548-3-PBS was coated to a plate as the capture antibody and incubated with serial dilutions of standard Eg2981. 87548-2-PBS was HRP conjugated as the detection antibody. Range: 0.156-10 ng/mL.