

For Research Use Only

Phospho-PAK4 (Ser474)/PAK5 (Ser602)/PAK6 (Ser560) Recombinant monoclonal antibody, PBS Only

Catalog Number: 87648-2-PBS



Basic Information

Catalog Number: 87648-2-PBS	GenBank Accession Number: NM_001014831	Purification Method: Protein A purification
Size: 100ug , Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 10298	CloneNo.: 253138E10
Source: Rabbit	UNIPROT ID: O96013	
Isotype: IgG	Full Name: p21 protein (Cdc42/Rac)-activated kinase 4	
	Calculated MW: 64 kDa	
	Observed MW: 72 kDa	

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
human, mouse

Background Information

PAK4(p21-activated kinase 4) is also named as KIAA1142. It belongs to the protein kinase superfamily. Phospho-PAK4 (Ser474)/PAK5 (Ser602)/PAK6 (Ser560) refers to the phosphorylated forms of the Group II p21-Activated Kinases (PAKs) at specific serine residues in their kinase activation loops. These phosphorylation events are essential for the catalytic activation of Group II PAKs and serve as definitive markers of their enzymatic activity. (PMID: 17292838)

Storage

Storage:
Store at -80°C.

Storage Buffer:
PBS only, pH7.3



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated HeLa cells, phosphatase inhibitor treated HeLa cells and λ phosphatase treated HeLa cells were subjected to SDS PAGE followed by western blot with 87648-2-RR (Phospho-PAK4 (Ser474)/PAK5 (Ser602)/PAK6 (Ser560) antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH (60004-1-Ig) antibody as a loading control. This data was developed using the



Non-treated NIH/3T3 cells, phosphatase inhibitor treated NIH/3T3 cells and λ phosphatase treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 87648-2-RR (Phospho-PAK4 (Ser474)/PAK5 (Ser602)/PAK6 (Ser560) antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH (60004-1-Ig) antibody as a loading control. This data was developed using the

