

For Research Use Only

NLGN3 Recombinant monoclonal antibody, PBS Only

Catalog Number: 87774-2-PBS



Basic Information

Catalog Number: 87774-2-PBS	GenBank Accession Number: NM_181303	Purification Method: Protein A purification
Size: 100ug, Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 54413	CloneNo.: 253199D2
Source: Rabbit	UNIPROT ID: Q9NZ94-1	
Isotype: IgG	Full Name: neuroligin 3	
Immunogen Catalog Number: EG5477	Calculated MW: 94 kDa	
	Observed MW: 90 kDa, 110 kDa	

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
human, mouse, rat

Background Information

Neuroligins (NLGNs) are cell adhesion molecules located on the postsynaptic side of the synapse that interact with their presynaptic partners neuroligins to maintain trans-synaptic connection. Three highly homologous neuroligins are abundantly expressed in rodent brain: NLGN1, NLGN2, NLGN3, while NLGN4 is less abundant in rodent brain and less evolutionarily conserved among all NLGNs isoforms. The literature indicates that significantly elevated basal levels of cross-linked NLGN3-positive dimers (sNLGN3, ~300 kDa) can be detected on the synaptic surface of ^{*}Fmr1[±] KO mice. And also Intracellular NLGN3 (iNLGN3 at about 120 kDa) level was increased in basal state Fmr1 KO synaptoneuroosomes (PMID: 30056576).

Storage

Storage:
Store at -80°C.

Storage Buffer:
PBS only, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 87774-2-RR (NLGN3 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 87774-2-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 87774-2-RR (NLGN3 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 87774-2-PBS in a different storage buffer formulation.