For Research Use Only

Anti-Mouse IL-12/IL-23 p40 Rabbit Recombinant Antibody, PBS Only

Proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Catalog Number: 98004-1-PBS

Basic Information

Catalog Number: 98004-1-PBS

Size: 1mg,2mg/ml Source: Rabbit GeneID (NCBI): 16160 UNIPROT ID: P43432 Full Name:

GenBank Accession Number:

interleukin 12b Calculated MW: 70 kDa Purification Method: Protein A purification

CloneNo.: 230037C9

Applications

Tested Applications:

FC (Intra)

Isotype:

Species Specificity:

mouse

Background Information

IL-12 and IL-23 are heterodimeric cytokines that share a common p40 subunit (PMID: 11114383). IL-12 is composed of the IL-12 p40 subunit linked to the IL-12 p35 subunit, and the heterodimer signals through the IL-12 receptor (IL-12R), which comprises the IL-12R β 1 and IL-12R β 2 subunits. IL-23 is composed of the IL-23 p19 subunit and the IL-12 p40 (IL-12/23p40) subunit, which signals through IL-23R and IL-12R β 1 (PMID: 11114383; 26121196). IL-12/IL-23 p40 also exists as a monomer and as a homodimer which can act as a potent IL-12 antagonist (PMID: 8958912; 18783467). IL-12/IL-23 p40 is produced by antigen-presenting cells, such as dendritic cells (DCs), monocytes, macrophages, neutrophils and, to a lesser extent, B cells (PMID: 20476918).

Storage


Storage:


Store at -80°C. Stable for one year after shipment.

Storage Buffer:

PBS only

Selected Validation Data





Biolayer interferometry (BLI) kinetic assays of 98004-1-RR against Mouse IL-12/23 p40 were performed. The affinity constant is 4.18 nM.

1X10^6 unstimulated or IFNy and LPS treated (in the presence of protein transport inhibitors) RAW 264.7 cells were intracellularly stained with 0.25 ug Anti-Mouse IL-12/23 p40 (98004-1-PB5, Clone: 230037C9) and PE-Conjugated Affini Pure Goat Anti-Rabbit IgG(H+L) at dilution 1:700. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

1X10^6 unstimulated or IFNy and LPS treated (in the presence of protein transport inhibitors) RAW 264.7 cells were intracellularly stained with 0.125 ug Anti-Mouse IL-12/IL-23 p40 (98004-1-RR, Clone: 230037C9) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:700. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 98004-1-