For Research Use Only

Anti-Human CD141/Thrombomodulin Rabbit Recombinant Antibody

Catalog Number: 98037-1-RR



Basic Information

Catalog Number:

98037-1-RR

Size:

100ug , 1 mg/ml

Source: Rabbit Isotype: GenBank Accession Number:

NM_000361.3 GeneID (NCBI):

7056

UNIPROT ID: P07204

Full Name: thrombomodulin

Calculated MW:

60 kDa

Purification Method: Protein A purfication

CloneNo.: 240365E11

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

Thrombomodulin, also known as CD141, is an endothelial cell surface glycoprotein that forms a 1:1 complex with the coagulation factor thrombin and plays an important role as a natural anticoagulant. Thrombomodulin serves to convert thrombin from a procoagulant protein into the activator for protein C. Once converted to activated protein C (APC), this protein serves as a major anticoagulant in blood (PMID: 2827310). Thrombomodulin is also located in other cells (keratinocytes, osteoblasts, macrophages,...) where it might be involved in cell differentiation or in inflammation (PMID: 9814688). In humans, thrombomodulin is encoded by the THBD gene. Mutations in this gene are a cause of thromboembolic disease, also known as inherited thrombophilia. Thrombomodulin is glycosylated and has an apparent molecular weight of 75 to 110 kDa (PMID: 1650405; 2827310).

Storage


Storage

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.


Selected Validation Data



1x10^6 human peripheral blood leukocytes were surface stained with 0.25 ug Anti-Human CD141/Thrombomodulin Rabbit Recombinant Antibody (98037-1-RR, Clone: 240365E11) or Isotype Control and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining.Cells were not fixed.



1x10^6 human peripheral blood leukocytes were surface stained with 0.25 ug Anti-Human Thrombomodulin Rabbit Recombinant Antibody (98037-1-RR, Clone: 240365E11) (red) or Isotype Control (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



Biolayer interferometry (BLL) kinetic assays of 98037-1-RR against Human CD141/Thrombomodulin were performed. The affinity constant is 1.83 nM.