For Research Use Only

Anti-Mouse ICOSLG/B7-H2/CD275 Rabbit Recombinant Antibody, PBS Only

Uni-rAb www.ptglab.com

Purification Method:

Protein A purfication

CloneNo.:

240243G5

Catalog Number:98058-1-PBS

Basic Information

Catalog Number:

98058-1-PBS

Size:

1mg, 2mg/ml Source:

Rabbit Isotype: GenBank Accession Number:

NM_015790.3 GeneID (NCBI):

50723

UNIPROT ID: Q9JHJ8-1

Full Name: icos ligand

Calculated MW: 36 kDa

Applications

Tested Applications:

Species Specificity:

mouse


Background Information

ICOSLG, also known as B7H2 or GL50, is a type I transmembrane glycoprotein belonging to the B7 ligand family. ICOSLG is extensively expressed on professional antigen-presenting cells including B cells, macrophages, and dendritic cells, as well as non-lymphoid cells including mesenchymal stem cells, endothelial cells, fibroblasts, and tumor cells (PMID: 33756276; 35800767). It is a specific ligand for the T-cell-specific cell surface receptor ICOS and $acts as a co-stimulatory \ molecule \ (PMID: 11023515; 11007762). \ The interaction of ICOSLG \ and \ ICOS \ plays \ important$ roles in the activation, proliferation, differentiation, and cytokine production of T cells as well as in the antibody secretion from B cells during secondary immune responses (PMID: 21851236).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data



1x10^6 mouse splenocytes were surface stained with 0.25 ug Anti-Mouse ICOSLG/B7-H2/CD275 Rabbit Recombinant Antibody (98058-1-RR, Clone:240243G5) or Isotype Control and PE-conjugated Goat Anti-Rabbit IgG. Cells were then stained with Coralite® Plus 647 Anti-Mouse CD45R (B220). Cells were not fixed. This data was developed using the same antibody clone with 98058-1-PBS in a different storage buffer formulation.



1x10^6 mouse splenocytes were surface stained with 0.25 ug Anti-Mouse ICOSLG/B7-H2/CD275 Rabbit Recombinant Antibody (98058-1-RR, Clone:240243G5) (red) or Isotype Control (blue) and PE-conjugated Goat Anti-Rabbit IgC. Cells were not fixed. This data was developed using the same antibody clone with 98058-1-PBS in a different storage buffer formulation.



Biolayer interferometry (BLL) kinetic assays of 98058-1-RR against Mouse ICOSLG/B7-H2/CD275 were performed. The affinity constant is below 1 pM.