For Research Use Only

## Anti-Mouse CD40L/CD154 Rabbit Recombinant Antibody, PBS Only

www.ptglab.com

**Purification Method:** 

Protein A purfication

CloneNo.:

240743E6

Catalog Number: 98120-1-PBS

**Basic Information** 

Catalog Number:

98120-1-PBS

Size:

1mg, 2 mg/ml Source: Rabbit

Isotype: IgG

GenBank Accession Number:

NM\_011616.2 GeneID (NCBI):

21947

**UNIPROT ID:** P27548 Full Name: CD40 ligand

Calculated MW: 29 kDa

**Applications** 

**Tested Applications:** 

Species Specificity:

mouse


**Background Information** 

The CD40 ligand (CD40L, TRAP, CD154), a member of the TNF superfamily of ligands, is expressed as either a 33-kd transmembrane homologue or 18-kd soluble form (sCD154). CD40L is primarily expressed on activated CD4+ T cells and on a small proportion of CD8+ T cells and platelets. It binds to CD40 on antigen-presenting cells (APC), which  $leads \ to \ many \ effects \ depending \ on \ the \ target \ cell \ type. \ It \ has \ been \ suggested \ that \ CD40/CD40L \ interactions$ regulate oxidative stress and affect various signaling pathways in both the immunological and the cardiovascular systems. The CD40/CD40L system is also involved in tumorigenesis. Its expression is tightly regulated, and the control of thabnormal levels of CD40L are associated with the pathogenesis of atheromatous plaque destabilization and thrombotic events. Multiple mutations in CD40LG gene have been identified that are associated with hyper-IgM immunodeficiency syndrome type 1.


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only

## **Selected Validation Data**



1x10^6 untreated or PMA and ionomycin treated mouse CD3+ T cells were surface stained with 0.25 ug Anti-Mouse CD40L/CD154 Rabbit Recombinant Antibody (98120-1-RR, Clone: 240743E6) and PEconjugated Goat Anti-Rabbit IgG. Cells were costained with Coralite® Plus 647 Anti-Mouse CD4. Cells were not fixed. This data was developed using the same antibody clone with 98120-1-PBS in a different storage buffer formulation.



1x10^6 PMA and ionomycin treated mouse CD3+T cells were surface stained with 0.25 ug Anti-Mouse CD40L/CD154 Rabbit Recombinant Antibody (98120-1-RR, Clone: 240743E6) (red) or Isotype Control (blue) and PE-conjugated Goat Anti-Rabbit IgG. 1x10^6 untreated mouse CD3+T cells were surface stained with 0.25 ug Anti-Mouse CD40L/CD154 Rabbit Recombinant Antibody (98120-1-RR, Clone: 240743E6) and PE-conjugated Goat Anti-Rabbit IgG (black,