For Research Use Only

Anti-Human CD30/TNFRSF8 Rabbit Recombinant Antibody

Catalog Number:98124-1-RR





| Basic Information | Catalog Number: 98124-1-RR | GenBank Accession Number: NM_001243 | Purification Method: Protein A purfication | |
|-----------------------|---|--|---|--|
| | Size: 100ug , 1000 µg/ml | GeneID (NCBI): 943 | CloneNo.: 241330A12 | |
| | Source: Rabbit | UNIPROT ID: P28908 | | |
| | Isotype: IgG | Full Name: tumor necrosis factor receptor superfamily, member 8 | | |
| | | Calculated MW: 64 kDa | | |
| Applications | Tested Applications: FC | | | |
| | Species Specificity: human | | | |
| Background Informatio | receptor superfamily. It is co segment, and an intracellula lymphocytes, and a variety c and anaplastic large cell lym transmembrane glycoproteir | CD30, also known as TNFRSF8 or Ki-1, is a transmembrane glycoprotein belonging to the tumor necrosis factor receptor superfamily. It is composed of an extracellular domain with six cysteine-rich repeats, a transmembrane segment, and an intracellular domain (PMID: 1310894). CD30 is expressed on a small subset of activated T and B lymphocytes, and a variety of lymphoid neoplasms, with the highest expression in classical Hodgkin's lymphoma and anaplastic large cell lymphoma (PMID: 9826579; 28885612). The CD30 ligand (CD30L or CD153) is a type II transmembrane glycoprotein of the tumor necrosis factor ligand superfamily (PMID: 8701986). Signal transduction of CD30/CD30L utilizes signal transducers, TNFR-associated factors (TRAF1, 2, 3 and 5) (PMID: 9826579). | | |
| Storage | Storage: Store at 2 - 8°C. Stable for one year after shipment. Storage Buffer: PBS with 0.09% sodium azide, pH 7.3. | | | |

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10^6 PHA-treated human PBMCs were surface stained with 0.25 ug Anti-Human CD30 Rabbit Recombinant Antibody (98124-1-RR, Clone: 241330A12) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (red), 0.25 ug or Isotype Control and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (blue). 1x10^6 untreated human PBMCs were surface stained with 0.25 ug Anti-Human CD30 Rabbit Recombinant Antibody (98124-1-RR, Clone: 241330A12)

Biolayer interferometry (BLI) kinetic assays of 98124-1-RR against Human CD30 were performed. The affinity constant is 0.528 nM.