For Research Use Only

Anti-Mouse CD27 Rabbit Recombinant Antibody

Catalog Number:98160-1-RR



Purification Method:

Protein A purfication

CloneNo.:

241378E11

Basic Information

Catalog Number:

98160-1-RR

Size:

100ug , 1000 $\mu g/ml$

Source: Rabbit Isotype: GenBank Accession Number:

NM_001033126.2

GeneID (NCBI):

21940

UNIPROT ID: P41272 Full Name:

CD27 antigen

Calculated MW: 28kDa

Applications

Tested Applications:

Species Specificity:

mouse

Background Information

CD27 (also known as TNFRSF7) is a type I glycoprotein expressed on some B cells and the majority of T cells, and is a member of the tumor necrosis factor (TNF) receptor family. CD27 is required for generation and long-term maintenance of T cell immunity (PMID: 11062504). It is a receptor for CD70 (CD27L). Ligation of CD27 by CD70 induces strong ubiquitination of TRAF and the activation of both canonical and non-canonical NF-kappaB pathways, as well as the JNK pathway (PMID: 19426224). CD27 may also play a role in apoptosis through association with SIVA1.


Storage


Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.09% sodium azide, pH 7.3.

Selected Validation Data






1x10^6 mouse splenocytes were surface stained with 0.25 ug Anti-Mouse CD27 Rabbit Recombinant Antibody (98160-1-RR, Clone:241378E11) or Isotype Control, and PE-Conjugated Goat Anti-Rabbit IgG(H+L). cells were co-stained with APC Anti-mouse CD3. Cells were not fixed.



1x10^6 mouse splenocytes were surface stained with 0.25 ug Anti-Mouse CD27 Rabbit Recombinant Antibody (98160-1-RR, Clone: 241378E11) (red), or Isotype Control (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed.



Biolayer interferometry (BLL) kinetic assays of 98160-1-RR against Mouse CD27 were performed. The affinity constant is 1.87 nM.