

For Research Use Only

Anti-Human CD98/SLC3A2 Rabbit Recombinant Antibody

Catalog Number: 98190-1-RR



Basic Information

Catalog Number:

98190-1-RR

Size:

100ug, 1000 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_002394.5

GeneID (NCBI):

6520

UNIPROT ID:

P08195-1

Full Name:

solute carrier family 3 (activators of dibasic and neutral amino acid transport), member 2

Calculated MW:

68kDa

Purification Method:

Protein A purification

CloneNo.:

241755D6

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

CD98 is a 125-kDa type II membrane glycoprotein heterodimer composed of a 40-kDa non-glycosylated light chain (amino-acid transporter, encoded by the SLC7A5) and an 85-kDa glycosylated heavy chain (CD98hc, encoded by the SLC3A2) (PMID: 11507094). CD98hc also interacts with certain integrin b-subunits to regulate cell migration, survival, proliferation, and adhesion/polarity. CD98hc is overexpressed on the cell surface of many cancers and increased CD98hc expression is associated with the development and progression of tumors (PMID: 25084765). Molecular orientation studies have demonstrated that the C-terminal tail of human CD98 (hCD98), which contains a PDZ-binding domain, is extracellular (PMID: 18625289).

Storage

Storage:

Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH 7.3.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human CD98 Rabbit Recombinant Antibody (98190-1-RR, Clone: 241755D6) or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human CD98 Rabbit Recombinant Antibody (98190-1-RR, Clone: 241755D6) (red) or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



Biolayer interferometry (BLI) kinetic assays of 98190-1-RR against Human CD98/SLC3A2 were performed. The affinity constant is 5.97 nM.