For Research Use Only

Anti-Mouse CD30/TNFRSF8 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98209-1-PBS



Purification Method:

Protein A purfication

CloneNo.:

241710B4

Basic Information

Catalog Number:

98209-1-PBS

Size:

1mg, 2 mg/ml Source: Rabbit

Isotype:

GenBank Accession Number:

NM_009401.2

GeneID (NCBI): 21941

UNIPROT ID:

Q60846 Full Name:

tumor necrosis factor receptor

superfamily, member 8

Calculated MW: 53kDa

Applications

Tested Applications:

Species Specificity:

mouse

Background Information

 ${\tt CD30, also \, known \, as \, TNFRSF8 \, or \, Ki-1, is \, a \, transmembrane \, glycoprotein \, belonging \, to \, the \, tumor \, necrosis \, factor \, is \, a \, transmembrane \, glycoprotein \, belonging \, to \, the \, tumor \, necrosis \, factor \, is \, a \, transmembrane \, glycoprotein \, belonging \, to \, the \, tumor \, necrosis \, factor \, is \, a \, transmembrane \, glycoprotein \, belonging \, to \, the \, tumor \, necrosis \, factor \, is \, a \, transmembrane \, glycoprotein \, belonging \, to \, the \, tumor \, necrosis \, factor \, is \, a \, transmembrane \, glycoprotein \, belonging \, to \, the \, tumor \, necrosis \, factor \, is \, a \, transmembrane \, glycoprotein \, belonging \, to \, the \, tumor \, necrosis \, factor \, is \, a \, transmembrane \, glycoprotein \, belonging \, to \, the \, tumor \, necrosis \, factor \, is \, a \, transmembrane \, glycoprotein \, belonging \, to \, the \, tumor \, necrosis \, factor \, is \, a \, transmembrane \, glycoprotein \, belonging \, to \, the \, tumor \, necrosis \, factor \, is \, a \, transmembrane \, glycoprotein \, belonging \, to \, the \, tumor \, necrosis \, factor \, is \, a \, transmembrane \, glycoprotein \, belonging \, to \, the \, tumor \, necrosis \, factor \, the \, tumor \, necrosis \, the \, tumor \, ne$ receptor superfamily. It is composed of an extracellular domain with six cysteine-rich repeats, a transmembrane segment, and an intracellular domain (PMID: 1310894). CD30 is expressed on a small subset of activated T and B lymphocytes, and a variety of lymphoid neoplasms, with the highest expression in classical Hodgkin's lymphoma and anaplastic large cell lymphoma (PMID: 9826579; 28885612). The CD30 ligand (CD30L or CD153) is a type II transmembrane glycoprotein of the tumor necrosis factor ligand superfamily (PMID: 8701986). Signal transduction of CD30/CD30L utilizes signal transducers, TNFR-associated factors (TRAF1, 2, 3 and 5) (PMID: 9826579).


Storage


Storage:

Store at -80°C. Storage Buffer:

PBS Only

Selected Validation Data





1x10^6 anti-CD3/CD28 treated mouse splenocytes were surface stained with 0.25 ug Anti-Mouse CD30/TNFRSF8 Rabbit Recombinant Antibody (98209-1-RR, Clone: 241710B4) (red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). 1x10^6 untreated mouse splenocytes were surface stained with 0.25 ug Anti-Mouse CD30/TNFRSF8 Rabbit Recombinant Antibody (98209-1-RR, Clone: 241710B4)



Biolayer interferometry (BLI) kinetic assays of 98209-1-RR against Mouse CD30/TNFRSF8 were performed. The affinity constant is 1.64 nM.