For Research Use Only

Anti-Mouse Ki-67 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98224-1-PBS



Purification Method:

Protein A purfication

CloneNo.:

241768E2

Basic Information

Catalog Number:

98224-1-PBS

Size:

1mg, 2 mg/ml Source:

Rabbit Isotype:

IgG

antigen identified by monoclonal antibody Ki 67

> Calculated MW: 351kDa

GenBank Accession Number:

NM_001081117.2

GeneID (NCBI):

UNIPROT ID: E9PVX6-1

17345

Applications

Tested Applications:

FC (Intra)

Species Specificity:

mouse

Background Information


The Ki-67 protein (also known as MKI67) is a cellular marker for proliferation. Ki67 is present during all active phases of the cell cycle (G1, S, G2 and M), but is absent in resting cells (G0). Cellular content of Ki-67 protein markedly increases during cell progression through S phase of the cell cycle. Therefore, the nuclear expression of Ki-67 can be evaluated to assess tumor proliferation by immunohistochemistry. It has been demonstrated to be of prognostic value in breast cancer. In head and neck cancer, several studies have reported an association between high proliferative activity and poorer prognosis.

Storage


Storage: Store at -80°C. Storage Buffer:

PBS Only

Selected Validation Data



1x10^6 untreated or Con A treated C57BL/6 mouse splenocytes were intracellularly stained with 0.25 ug Anti-Mouse Ki-67 Rabbit Recombinant Antibody (98224-1-RR, Clone: 241768E2) and PE-Conjugated Goat Anti-Rabbit 1gG(H+L). Cells were co-stained with CoraLite® Plus 647 Anti-Mouse CD3. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 98224-1-PBS



1x10^6 Con A treated C57BL/6 mouse splenocytes were intracellularly stained with 0.25 ug Anti-Mouse Ki-67 Rabbit Recombinant Antibody (98224-1-RR, Clone: 241768E2) (red) or 0.25 ug Rabbit 1gG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue) and PE-Conjugated Goat Anti-Rabbit 1gG(H+L). 1x10^6 untreated C57BL/6 mouse splenocytes were intracellularly stained with 0.25 ug Anti-Mouse Ki-67 Rabbit Recombinant Antibody

