## For Research Use Only

## Anti-Human CD46 Rabbit Recombinant Antibody, PBS Only

Catalog Number:98240-1-PBS




## Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 98240-1-PBS NM\_172351.3 Protein A purfication GeneID (NCBI): CloneNo.: Size: 1mg , 2 mg/ml 4179 241764C9 Source: UNIPROT ID: Rabbit P15529-11 Full Name: Isotype: lgG CD46 molecule, complement regulatory protein Immunogen Catalog Number: EG1770 Calculated MW: 42kDa **Applications Tested Applications:** FC Species Specificity: human **Background Information** CD46, also known as Membrane cofactor protein (MCP) or MIC10, is a ubiquitously expressed type type-I transmembrane glycoprotein that acts as a cofactor for complement factor I to mediate inactivation of C3b and C4b deposited on host cells. CD46 is a receptor for a large number of pathogens, including measles virus and adenovirus (PMID: 20941397). CD46 plays a role in linking innate and adaptive immune responses. It may be involved in the fusion of the spermatozoa with the oocyte during fertilization. Mutations of the CD46 gene have been associated with some diseases including hemolytic uremic syndrome (PMID: 26054645). CD46 is expressed on most cells as four isoforms that are derived via alternative splicing (PMID: 15324743). Storage: Storage Store at -80°C. Storage Buffer: PBS only, pH7.3


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1x10<sup>^6</sup> human PBMCs were surface stained with 0.125 ug Anti-Human CD46 Rabbit Recombinant Antibody (98240-1-RR, Clone: 241764C9) or Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were then stained with CoraLite® Plus 647 Anti-Human CD19. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same



1x10<sup>^6</sup> human PBMCs were surface stained with 0.125 ug Anti-Human CD46 Rabbit Recombinant Antibody (98240-1-RR, Clone: 241764C9) (red) or Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98240-1-PBS in a different storage