For Research Use Only

Anti-Human EMR2/CD312 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98260-1-PBS



Purification Method:

Protein A purfication

CloneNo.:

242042B11

Basic Information

Catalog Number:

98260-1-PBS

Size:

1mg, 2 mg/ml Source:

Rabbit Isotype:

IgG

GenBank Accession Number:

NM_013447.3

GeneID (NCBI):

30817 **UNIPROT ID:**

Q9UHX3-1

egf-like module containing, mucin-

like, hormone receptor-like 2

Calculated MW:

90 kDa

Applications

Tested Applications:

Species Specificity:

human

Background Information


EMR2 (Epidermal Growth Factor-like module-containing Mucin-like Hormone Receptor 2), also known as CD312 or ADGRE2, is a member of the adhesion-GPCR family that contains extracellular EGF-like domains. EMR2 contains a $total\ of\ five\ tandem\ EGF-like\ domains\ and\ expresses\ similar\ protein\ is oforms\ consisting\ of\ various\ numbers\ of\ EGF-like\ domains\ and\ expresses\ similar\ protein\ is\ of\ various\ numbers\ of\ EGF-like\ domains\ and\ expresses\ similar\ protein\ is\ of\ various\ numbers\ of\ EGF-like\ domains\ and\ expresses\ similar\ protein\ is\ of\ various\ numbers\ of\ expresses\ similar\ protein\ is\ of\ various\ numbers\ of\ expresses\ similar\ protein\ is\ of\ various\ numbers\ of\ expresses\ of\ expresses\ of\ expresses\ numbers\ of\ expresses\ of\ expre$ like domains as a result of alternative RNA splicing (PMID: 10903844). EMR2 is restricted to myeloid cells including monocytes, macrophages, dendritic cells and granulocytes28 and its expression is highly regulated during monocyte/macrophage differentiation (PMID: 31969668). The ligation of EMR2 could increase neutrophil adhesion, migration and anti-microbial mediator production and could enhance the systemic inflammation (PMID: 27905560).

Storage


Storage: Store at -80°C. Storage Buffer:

PBS Only


Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.125 ug Anti-Human EMR2/CD312
Rabbit Recombinant Antibody (98260-1-RR, Clone: 242042B11) or 0.125 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were co-stained with APC Anti-Human CD14. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same



1x10^6 human PBMCs were surface stained with 0.125 ug Anti-Human EMR2/CD312 Rabbit Recombinant Antibody (98260-1-RR, Clone: 242042B11)(red) or 0.125 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98260-1-PBS in a



Biolayer interferometry (BLI) kinetic assays of 98260-1-RR against Human EMR2/CD312 were performed. The affinity constant is 0.529 nM.