Anti-Mouse CD19 Rabbit Recombinant Antibody




Catalog Number:98287-1-RR


Basic Information	Catalog Number: 98287-1-RR	GenBank Accession Number: NM_009844.2	Purification Method: Protein A purfication
	Size: 100ug , 1000 µg/ml	GenelD (NCBI): 12478	CloneNo.: 242301C6
	Source: Rabbit	UNIPROT ID: P25918	
	Isotype: IgG	Full Name: CD19 antigen	
		Calculated MW: 60 kDa	
Applications	Tested Applications: FC		
	Species Specificity: mouse		
Background Information	CD19 is a 95 kDa type I transmembrane glycoprotein belonging to the immunoglobulin superfamily (PMID: 2472450). It is expressed by B cells and follicular dendritic cells. CD19 is up-regulated at the step of B-lineage commitment during the differentiation of the hematopoietic stem cell, it remains on during subsequent stages of differentiation until finally down-regulated during terminal differentiation into plasma cells (PMID: 8528044). CD19 is involved in B cell development, activation and differentiation. It is the dominant component for the signaling complex on B cells that includes CD21 (CR2), CD81 (TAPA-1) and CD225 and acts as a critical co-receptor for BCR signal transduction (PMID: 23210908).		
Storage	Storage: Store at 2 - 8°C. Stable for one Storage Buffer: PBS with 0.09% sodium azide		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10^6 mouse splenocytes were surface stained with 0.25 ug Anti-Mouse CD19 Rabbit RecAb (98287-1-RR, Clone:242301C6) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were then stained with APC Anti-Mouse CD3 (17A2) (APC-65077, Clone: 17A2). Cells were not fixed.

1x10^{^6} mouse splenocytes were surface stained with 0.25 ug Anti-Mouse CD19 Rabbit RecAb (98287-1-RR, Clone: 242301C6) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed.