For Research Use Only

## Anti-Human BST2 Rabbit Recombinant Antibody

Catalog Number: 98305-2-RR



**Purification Method:** 

Protein A purification

Recommended Dilutions:

CloneNo.:

250317H5

**Basic Information** 

Catalog Number: GenBank Accession Number: 98305-2-RR

BC033873

GeneID (NCBI): 100ug , 1000  $\mu$ g/ml

**UNIPROT ID:** Source: Rabbit Q10589

FC: 0.25 ug per 10^6 cells in a 100  $\mu$ l suspension Full Name: Isotype:

bone marrow stromal cell antigen 2

Calculated MW: Immunogen Catalog Number: 180 aa, 20 kDa

**Applications** 

**Tested Applications:** 

Species Specificity:

human

Positive Controls: FC: human PBMCs,

## **Background Information**

BST2, also named as CD317 and Tetherin, belongs to the tetherin family. It may be involved in the sorting of secreted proteins and it is involved in pre-B-cell growth. BST2 is an antiretroviral defense protein, that blocks release of retrovirus from the cell surface. Depleted unpon HIV-1 infection by viral VPU protein through 20S proteasome degradation. Depleted upon infection by human Kaposi's sarcoma-associated herpesvirus (KSHV)  $through\ ubiquitination\ and\ subsequent\ degradation.\ BST2\ may\ play\ a\ role\ in\ B-cell\ activation\ in\ rheumatoid$ arthritis. It is recently identified interferon-induced cellular proteins that restrict infections by retroviruses and filoviruses and of influenza virus and flaviviruses, respectively. BST2 is a plasma membrane proteins, tetherin inhibits virion particle release from infected cells. BST2 is effective against retroviruses and flavoviruses whilst IFITMs disrupt influenza and flavivirus infection. Observed MW of BST2 is 30-36 kDa (PMID: 19196977; 21237475).

Storage


Storage:

Store at 2 - 8°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.09% sodium azide, pH7.3

## **Selected Validation Data**



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human BST2 Rabbit RecAb (98305-2-RR, Clone: 250317H5) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were then stained with Coralite® Plus 647 Anti-Human CD14. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human BST2 Rabbit RecAb (98305-2-RR, Clone: 250317H5) (red) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.