For Research Use Only

Anti-Human CEACAM6/CD66c Rabbit Recombinant Antibody, PBS Only

Catalog Number:98329-3-PBS



Basic Information	Catalog Number: 98329-3-PBS	GenBank Accession Number: BC005008	Purification Method: Protein A purification
	Size:	GenelD (NCBI):	CloneNo.:
	1mg , 2 mg/ml	4680	241997G1
	Source:	UNIPROT ID:	
	Rabbit	P40199	
	Isotype:	Full Name:	
	IgG	carcinoembryonic antigen-relate cell adhesion molecule 6 (non-	d
		specific cross reacting antigen)	
		Calculated MW:	
		37 kDa	
Applications	Tested Applications: FC		
	Species Specificity:		
	human		
Background Information	Carcinoembryonic antigen-related cell adhesion molecules (CEACAMs) belong to the immunoglobulin superfamily of cell adhesion molecules. CEACAMs play major roles in cell-cell and cell-ECM adhesion and have been implicated in controlling cell proliferation, angiogenesis, and tissue remodeling (PMID: 23021083). CEACAM6, also known as CD66c, is a glycosylphosphatidylinositol (GPI)-linked cell surface protein that belongs to the CEACAM family of the immunoglobulin superfamily (PMID: 23903773). It is normally expressed on the surface of myeloid and epithelial surfaces and upregulated preferentially at the apical/luminal membranes of many tumors (PMID: 27595061; 35082925). CEACAM6 plays an important role in cell adhesion, proliferation, tumorigenesis, metastasis, and immunity (PMID: 31797958).		
Storage	Storage: Store at -80°C. Storage Buffer: PBS Only		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^{^6} human peripheral blood leukocytes were surface stained with 0.25 ug Anti-Human CEACAM6/CD66c Rabbit RecAb (98329-3-RR, Clone: 241997G1) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9), and PE-conjugated Goat Anti-Rabbit IgG. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98329-3-PBS in a different storage buffer formulation.



1x10^{^6} human peripheral blood leukocytes were surface stained with 0.25 ug Anti-Human CEACAM6/CD66c Rabbit RecAb (98329-3-RR, Clone:241997G1) (red) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PEconjugated Goat Anti-Rabbit IgG. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98329-3-PBS in a different storage