For Research Use Only

Anti-Human CSF1R/CD115 Rabbit Recombinant Antibody, PBS Only

Catalog Number:98379-3-PBS




Catalog Number: 98379-3-PBS	GenBank Accession Number: BC047521	Purification Method: Protein A purification
Size: 1mg, 2 mg/ml	GenelD (NCBI): 1436	CloneNo.: 250001D7
Source: Rabbit	UNIPROT ID: P07333	
Isotype: IgG	Full Name: colony stimulating factor 1 recep	tor
	Calculated MW: 972 aa, 108 kDa	
Tested Applications: FC		
Species Specificity: human		
The CSF-1 receptor (CSF1R), also called CD115, is activated by the homodimeric growth factors colony-stimulating factor-1 (CSF-1) and interleukin-34 (IL-34). It plays important roles in development and in innate immunity by regulating the development of most tissue macrophages and osteoclasts, of Langerhans cells of the skin, of Paneth cells of the small intestine, and of brain microglia. It also regulates the differentiation of neural progenitor cells and controls functions of oocytes and trophoblastic cells in the female reproductive tract. Owing to this broad tissue expression pattern, it plays a central role in neoplastic, inflammatory, and neurological diseases.		
Storage: Store at -80°C. Storage Buffer: PBS only, pH7.3		
	98379-3-PBS Size: 1mg, 2 mg/ml Source: Rabbit Isotype: IgG Tested Applications: FC Species Specificity: human The CSF-1 receptor (CSF1R) factor-1 (CSF-1) and interley regulating the developmen cells of the small intestine, controls functions of oocyte expression pattern, it plays Storage: Storage Buffer:	98379-3-PBS BC047521 Size: GeneID (NCBI): 1mg, 2 mg/ml 1436 Source: UNIPROT ID: Rabbit P07333 Isotype: Full Name: IgG colony stimulating factor 1 receptor Calculated MW: 972 aa, 108 kDa Tested Applications: FC Species Specificity: human The CSF-1 receptor (CSF1R), also called CD115, is activated by the hom factor-1 (CSF-1) and interleukin-34 (IL-34). It plays important roles in doregulating the development of most tissue macrophages and osteoclass cells of the small intestine, and of brain microglia. It also regulates the controls functions of oocytes and trophoblastic cells in the female reprocexpression pattern, it plays a central role in neoplastic, inflammatory, and storage: Storage: Storage Buffer:


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CSF1R/CD115 Rabbit RecAb (98379-3-RR, Clone:250001D7) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98379-3-PBS in a different storage buffer formulation.



1x10^{^6} human PBMCs were surface stained with 0.25 ug Anti-Human CSF1R/CD115 Rabbit RecAb (98379-3-RR, Clone: 250001D7) (red) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98379-3-PBS in a different storage buffer formulation.