For Research Use Only

Anti-Mouse OX40L/TNFSF4 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98381-1-PBS



Purification Method:

Protein A purification

CloneNo.:

242227B1

Basic Information

Catalog Number:

98381-1-PBS

Size:

1mg, 2 mg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

GenBank Accession Number:

NM_009452.2

GeneID (NCBI): 22164

UNIPROT ID: P43488

Full Name: tumor necrosis factor (ligand)

superfamily, member 4

Calculated MW: 22kDa

Applications

Tested Applications:

Species Specificity:

mouse

Background Information


The tumour necrosis factor ligand superfamily member 4 gene (TNFSF4, OX40L), which encodes for the T cell costimulatory molecule OX40 ligand, has been identified as a susceptibility gene for the development of systemic lupus erythematosus (SLE). OX40L/TNFSF4/CD252 is a co-stimulatory checkpoint protein expressed by several types of immune and non-immune cells (PMID: 15750594, 19778912). TNFSF4 mRNA is expressed in melanoma cell lines and melanoma samples, including those with low lymphocytic infiltrates, and is not associated with the ulceration status of the primary tumor (PMID: 31501955).

Storage


Storage: Store at -80°C. Storage Buffer:

PBS only, pH7.3

Selected Validation Data



1x10^6 untreated or Anti-IgM and CD40 treated BALB/c mouse splenocytes were surface stained with 0.25 ug Anti-Mouse 0X40 Ligand/TNFSF4 Rabbit RecAb (98381-1-RR, Clone:242227B1) and PE-Conjugated Goat Anti-Rabbit IgC(H+L). Cells were then stained with CoraLite® Plus 488 Anti-Mouse CD19 (1D3) (CL488-65290, Clone: 1D3). Cells were not fixed. This data was developed using the same antibody clone with 98381-1-PBS in a different storage buffer formulation.



1x10^6 Anti-IgM and CD40 treated BALB/c mouse splenocytes were surface stained with 0.25 ug Anti-Mouse 0X40 Ligand/TNFSF4 Rabbit RecAb (98381-1-RR, Clone:242227B1) (red) and PE-Conjugated Goat Anti-Rabbit IgG(H+L), or Rabbit IgG (Sotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). 1x10^6 untreated mouse splenocytes were surface stained with 0.25 ug Anti-Mouse 0X40 Ligand/TNFSF4 Rabbit RecAb (98381-1-RR, Clone:242227B1) (black,

