For Research Use Only

Anti-Human CD96 Rabbit Recombinant Antibody

Catalog Number: 98386-3-RR



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

98386-3-RR

NM_198196

Protein A purification

Size:

GeneID (NCBI): 10225 CloneNo.: 242964G12

100ug , 1000 μg/ml Source:

UNIPROT ID:

Recommended Dilutions:

Rabbit Isotype:

IgG

P40200 Full Name:

FC: 0.25 ug per 10^6 cells in 100 μ l suspension

CD96 molecule Calculated MW:

402 aa, 46 kDa

Applications

Tested Applications:

FC

Positive Controls:

Species Specificity:

human

FC: human PBMCs,

Background Information

CD96 is a member of the poliovirus receptor (PVR, CD155)-nectin family that includes T cell Ig and ITIM domain (TIGIT) and CD226. CD96 expression is broadly similar between mice and humans, and is present on a proportion of hematopoietic stem cells, $\alpha\beta$ and $\gamma\delta$ T-cells, NK cells and a sub-population of B-cells in humans and present on $\alpha\beta$ and $\gamma\delta$ T-cells, NK cells in mice (PMID: 32043568, 29531070)

Storage


Storage:

Store at 2 - 8°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.09% sodium azide, pH7.3

Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD96 Rabbit RecAb (98386-3-RR, Clone:242964G12), or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were then stained with CoraLite® Plus 647 Anti-Human CD19. Cells were not fixed. Lymphocyte were gated.



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD96 Rabbit RecAb (98386-3-RR, Clone:242964G12)(red), or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed. Lymphocyte were gated.