

For Research Use Only

Anti-Mouse CCL11/Eotaxin Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98464-1-PBS



Basic Information

Catalog Number:

98464-1-PBS

Size:

1mg, 3 mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG2400

GenBank Accession Number:

NM_011330

GeneID (NCBI):

20292

UNIPROT ID:

P48298

Full Name:

chemokine (C-C motif) ligand 11

Calculated MW:

11kd

Purification Method:

Protein A purification

CloneNo.:

243012F8

Applications

Tested Applications:

FC (Intra)

Species Specificity:

mouse

Background Information

CCL11 also named Eotaxin-1, was initially discovered as an eosinophil-specific chemoattractant. CCL11 plays a central role in both allergic and non-allergic inflammatory reactions by recruiting immune cells such as eosinophils, basophils and Th2 lymphocytes. CCL11 binds to the chemokine receptors CCR2, CCR3 and CCR5, with highest affinity to CCR3. High levels of CCL11 have been described in several chronic inflammatory diseases, such as allergic rhinitis, atopic dermatitis, asthma, gastrointestinal disease and rheumatoid arthritis.

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS only, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ HEK-293T cells (left) or CCL11/Eotaxin-transfected HEK-293T cells (right) were intracellularly stained with 0.25 ug Anti-Mouse CCL11/Eotaxin Rabbit RecAb (98464-1-RR, Clone: 243012F8) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 98464-1-PBS in a different storage buffer formulation.

1x10⁶ HEK-293T cells (left) or CCL11/Eotaxin-transfected HEK-293T cells (right) were intracellularly stained with 0.25 ug Anti-Mouse CCL11/Eotaxin Rabbit RecAb (98464-1-RR, Clone: 243012F8) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same

