

For Research Use Only

Anti-Mouse CCL9 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98475-1-PBS



Basic Information

Catalog Number:	98475-1-PBS	GenBank Accession Number:	NM_011338.2	Purification Method:	Protein A purification
Size:	1mg, 3 mg/ml	GeneID (NCBI):	20308	CloneNo.:	243103H5
Source:	Rabbit	UNIPROT ID:	Q3U9T8		
Isotype:	IgG	Full Name:	chemokine (C-C motif) ligand 9		
Immunogen Catalog Number:	EG3336	Calculated MW:	14KD		

Applications

Tested Applications:
FC (Intra)

Species Specificity:
mouse

Background Information

Mouse C-C motif ligand 9 (CCL9), alternatively named macrophage inflammatory protein 1γ (MIP-1γ), was identified in 1995 and is homologous to the mouse CCL6, as well as human CCL23 and CCL15. It is noted that CCL9 is also known by various names such as macrophage inflammatory protein-1 gamma (MIP-1γ), macrophage inflammatory protein-related protein-2 (MRP-2) and CCF18 in rodents. Monocytes and myeloid cell lines produce large quantities of CCL9, as do dendritic cells and T cells, in particular Th1 type T cells. Despite high baseline levels in circulation, it has become apparent that concentrations of CCL9 vary greatly in specific tissues with profound effects on health. For example, in the bone, CCL9 is produced at even higher levels, and is critical to osteoclast versus osteoblast differentiation of macrophages. There are also indications of a timed specific induction of CCL9 in skin wound healing and follicle-associated epithelium of the gut.


Storage

Storage:
Store at -80°C.
Storage Buffer:
PBS only, pH7.3

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ HEK-293T cells (left) or CCL9-transfected HEK-293T cells (right) were intracellularly stained with 0.25 ug Anti-Mouse CCL9 Rabbit RecAb (98475-1-RR, Clone: 243103H5) and APC-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 98475-1-PBS in a different storage buffer formulation.

1x10⁶ HEK-293T cells (left) or CCL9-transfected HEK-293T cells (right) were intracellularly stained with 0.25 ug Anti-Mouse CCL9 Rabbit RecAb (98475-1-RR, Clone:243103H5) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and APC-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same

