For Research Use Only

Anti-Rat CD8a Rabbit Recombinant Antibody Catalog Number:98484-2-RR



Basic Information	Catalog Number: 98484-2-RR	GenBank Accession Number: NM_031538.2	Purification Method: Protein A purification
	Size: 100ug , 1000 µg/ml	Genel D (NCBI): 24930	CloneNo.: 250721G1
	Source: Rabbit Isotype:	UNIPROT ID: P07725 Full Name:	Recommended Dilutions: FC: 0.25 ug per 10^6 cells in a 100 µl suspension
	IgG Immunogen Catalog Number: EG3839	CD8a molecule Calculated MW: 26 kDa	
Applications	Tested Applications: FC Species Specificity: rat	Positive Controls: FC : rat splenocytes,	
Background Information	CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8a β heterodimer while CD8aa homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) $\gamma \delta$ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).		
Storage	Storage: Store at 2 - 8°C. Stable for one yea Storage Buffer: PBS with 0.09% sodium azide, pH		

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10^6 rat splenocytes were surface stained with 0.25 ug Anti-Rat CD8a Rabbit RecAb (98484-2-RR, Clone:250721G1) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were then stained with FITC Plus Anti-Rat CD4 (OX-35) (FITC-65179, Clone: OX-35). Cells were not fixed.

1x10^6 rat splenocytes were surface stained with 0.25 ug Anti-Rat CD8a Rabbit RecAb (98484-2-RR, Clone: 250721G1) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (bue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed.