

For Research Use Only

# Anti-Human CXCL7/PPBP Rabbit Recombinant Antibody

Catalog Number: 98578-2-RR



## Basic Information

Catalog Number:

98578-2-RR

Size:

100ug, 1000 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG3690

GenBank Accession Number:

NM\_002704.3

GeneID (NCBI):

5473

UNIPROT ID:

P02775

Full Name:

pro-platelet basic protein (chemokine (C-X-C motif) ligand 7)

Calculated MW:

14kDa

Purification Method:

Protein A purification

CloneNo.:

250712D2

Recommended Dilutions:

FC (Intra): 0.25 ug per 10<sup>6</sup> cells in a 100 µl suspension

## Applications

Tested Applications:

FC (Intra)

Species Specificity:

human

Positive Controls:

FC (Intra) : Transfected CHO,

## Background Information

PPBP, also named as NAP2, or  $\beta$ -Thromboglobulin, is a platelet-derived growth factor that belongs to the CXC chemokine family. This growth factor is a potent chemoattractant and activator of neutrophils. It has been shown to stimulate various cellular processes including DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E2 secretion, and synthesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by synovial cells.

## Storage

Storage:

Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3





For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



1x10<sup>6</sup> CHO cells or CXCL7/PPBP transfected CHO were intracellularly stained with 0.25 ug Anti-Human CXCL7/PPBP Rabbit RecAb (98578-2-RR, Clone:250712D2) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00017).

1x10<sup>6</sup> CHO cells or CXCL7/PPBP transfected CHO were intracellularly stained with 0.25 ug Anti-Human CXCL7/PPBP Rabbit RecAb (98578-2-RR, Clone:250712D2)(red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00017).