

For Research Use Only

Anti-Human PTPRJ/CD148 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98715-1-PBS



Basic Information

Catalog Number: 98715-1-PBS	GenBank Accession Number: NM_002843.4	Purification Method: Protein A purification
Size: 1mg, 3 mg/ml	GeneID (NCBI): 5795	CloneNo.: 252382G6
Source: Rabbit	UNIPROT ID: Q12913-1	
Isotype: IgG	Full Name: protein tyrosine phosphatase, receptor type, J	
Immunogen Catalog Number: EG3492	Calculated MW: 146 kDa	

Applications

Tested Applications:
IHC, FC

Species Specificity:
human

Background Information

PTPRJ, also named as CD148 and DEP1, belongs to the protein-tyrosine phosphatase family and Receptor class 3 subfamily. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. PTPRJ may contribute to the mechanism of contact inhibition of cell growth. PTPRJ is present in all hematopoietic lineages, and was shown to negatively regulate T cell receptor signaling possibly through interfering with the phosphorylation of Phospholipase C Gamma 1 (PLCG1) and Linker for Activation of T Cells (LAT). It was also found to dephosphorylate PDGF beta receptor, and may be involved in UV-induced signal transduction.

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS only, pH7.3


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




1x10⁶ human peripheral blood lymphocytes were surface stained with 0.25 ug Anti-Human PTPRJ/CD148 Rabbit RecAb (98715-1-RR, Clone: 252382G6) or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98715-1-PBS in a



1x10⁶ human peripheral blood lymphocytes were surface stained with 0.25 ug Anti-Human PTPRJ/CD148 Rabbit RecAb (98715-1-RR, Clone: 252382G6) or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98715-1-PBS in a



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 98715-1-RR (DEP-1/CD148 antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 98715-1-PBS in a different storage buffer formulation.



Bi-layer interferometry (BLI) kinetic assays of 98715-1-RR against Human PTPRJ/CD148 were performed. The affinity constant is 0.649 nM.