

For Research Use Only

# Anti-Human HLA class II Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98748-1-PBS



## Basic Information

Catalog Number:

98748-1-PBS

Size:

1mg, 3 mg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

GeneID (NCBI):

Full Name:

Purification Method:

Protein A purification

CloneNo.:

253538A1

## Applications

Tested Applications:

FC

Species Specificity:

human

## Background Information

Human major histocompatibility complex (MHC) antigens, also referred to as human leukocyte antigens (HLA), are encoded by genes located on the short arm of chromosome 6 (6p21.3). There are two classes of HLA antigens: class I (HLA-A, B and C) and class II (HLA-D). This class I molecules are polymorphic membrane glycoproteins composed of a heavy (alpha) chain (44 kDa) which is encoded by a HLA class I gene (HLA-A, B or C), and  $\beta$ 2-microglobulin light (beta) chain (12 kDa). They are involved in the presentation of foreign antigens to the immune system (PMID: 667938; 3375250). This antibody can recognize HLA-DR, -DP, and -DQ.

## Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS only, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1x10<sup>6</sup> human PBMCs were surface stained with 0.25 ug Anti-Human HLA class II Rabbit RecAb (98748-1-RR, Clone:253538A1) and PE-Conjugated Goat Anti-Rabbit IgG(H+L), and 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9). Cells were then stained with CoraLite® Plus 647 Anti-Human CD14 Rabbit Recombinant Antibody (CL647-98040, Clone: 230332D7). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was



1x10<sup>6</sup> human PBMCs were surface stained with 0.25 ug Anti-Human HLA class II Rabbit RecAb (98748-1-RR, Clone:253538A1) and PE-Conjugated Goat Anti-Rabbit IgG(H+L), and 0.25 ug Anti-Human HLA-DR (L243) Mouse IgG2a Recombinant Antibody (65560-1-MR, Clone: L243). Cells were then stained with CoraLite® Plus 647 Anti-Human CD14 Rabbit Recombinant Antibody (CL647-98040, Clone: 230332D7). Cells were incubated with FC Receptor Block prior to staining. Cells

