

For Research Use Only

Anti-Human Integrin alpha-9 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98914-1-PBS



Basic Information

Catalog Number:

98914-1-PBS

Size:

1mg, 3 mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG6174

GenBank Accession Number:

NM_002207.2

GeneID (NCBI):

3680

UNIPROT ID:

Q13797

Full Name:

integrin, alpha 9

Calculated MW:

114 kDa

Purification Method:

Protein A purification

CloneNo.:

260170G5

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated α and β subunits (PMID: 9779984). Integrin alpha 9 contains a large N-terminal extracellular domain with 7 conserved repeats of putative metal-binding domains, a transmembrane segment, and a short C-terminal cytoplasmic tail (PMID: 8245132). Integrin alpha 9 interacts with integrin beta 1 generating $\alpha 9 \beta 1$ heterodimer, which is expressed in many cell types including epithelial cells, neutrophils, hepatocytes, muscle and endothelial cells (PMID: 21975548). It binds to a diversity of ligands in the extracellular matrix, like the α -disintegrin and metalloprotease (ADAM) family, THBS1, EMILIN1, VEGF, tenascin-C, osteopontin, fibronectin, and VCAM1 (PMID: 21975548; 33790909). Integrin $\alpha 9 \beta 1$ has been implicated in diverse biological functions including angiogenesis, lymphangiogenesis, lymphatic valve morphogenesis, granulopoiesis and tumorigenesis (PMID: 20179422).

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS only, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ human peripheral blood leukocytes were surface stained with 0.25 ug Anti-Human Integrin alpha-9 Rabbit RecAb (98914-1-RR, Clone: 260170G5) or Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were then stained with FcZero-rAb® APC-Cyanine7 Anti-Human CD15 Rabbit Recombinant Antibody (AY7-FcA98616, Clone: 252342A1). Cells were incubated with



1x10⁶ human peripheral blood leukocytes were surface stained with 0.25 ug Anti-Human Integrin alpha-9 Rabbit RecAb (98914-1-RR, Clone: 260170G5) (red) or Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98914-1-PBS in a