

For Research Use Only

# Atlantic Blue™ Anti-Mouse CD8a (53-6.7)



Catalog Number: **AB-65069**

## Basic Information

|                                    |   |   |
|------------------------------------|---|---|
| <b>Catalog Number:</b><br>AB-65069 | <b>GenBank Accession Number:</b><br>BC030679  | <b>Purification Method:</b><br>Affinity purification              |
| <b>Size:</b><br>100ug, 0.5 mg/ml   | <b>GeneID (NCBI):</b><br>12525                | <b>CloneNo.:</b><br>53-6.7  |
| <b>Source:</b><br>Rat              | <b>Full Name:</b><br>CD8 antigen, alpha chain | <b>Excitation/Emission maxima wavelengths:</b><br>404 nm / 458 nm |
| <b>Isotype:</b><br>IgG2a, kappa    |   |   |

## Applications

**Tested Applications:**  
FC

**Species Specificity:**  
Mouse

## Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8αβ heterodimer while CD8αα homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) γδ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).

## Storage


**Storage:**  
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

**Storage Buffer:**  
PBS with 0.09% sodium azide.

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)      E: proteintech@ptglab.com  
W: ptglab.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



1x10<sup>6</sup> mouse splenocytes were surface co-stained with Anti-Mouse CD4 and 0.5 ug Atlantic Blue™ Anti-Mouse CD8a (AB-65069, Clone: 53-6.7) or 0.5 ug isotype control. Cells were not fixed.