For Research Use Only

## Atlantic Blue™ Anti-Mouse CD8a (53- proteintech 6.7)



Catalog Number: AB-65069

**Basic Information** 

Catalog Number:

GenBank Accession Number:

CD8 antigen, alpha chain

**Purification Method:** 

AB-65069 Size:

GeneID (NCBI):

Affinity purification CloneNo.:

100ug , 0.5 mg/ml

12525

BC030679

53-6.7

Source:

**UNIPROT ID:** P01731

Excitation/Emission maxima wavelengths:

Isotype:

Full Name:

404 nm / 458 nm

IgG2a, kappa

**Applications** 

**Tested Applications:** 

Species Specificity:

Mouse

**Background Information** 

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8αβ heterodimer while CD8αα homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR)  $\gamma\delta$  T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide.

in USA), or 1(312) 455-8498 (outside USA)

## Selected Validation Data



1x10^6 mouse splenocytes were surface stained with 0.5 ug Atlantic Blue™ Anti-Mouse CD8a (53-6.7) (AB-65069, Clone:53-6.7) or 0.5 ug Isotype Control. Cells were co-stained with 0.5 ug CoraLite® Plus 647 Anti-Mouse CD4 (GK1.5) (CL647-65104, Clone: GK1.5). Cells were not fixed.