For Research Use Only

Atlantic Blue™ Anti-Human CD25/IL-2RA (BC96)

Catalog Number: AB-65096



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Affinity purification

AB-65096 Size:

GeneID (NCBI):

CloneNo.:

100 tests , 5 μ l/test

3559

BC156627

BC96

Source: Mouse

UNIPROT ID: P01589

Excitation/Emission maxima wavelengths:

Isotype: lgG1, kappa Full Name: interleukin 2 receptor, alpha 404 nm / 458 nm

Calculated MW: 272 aa, 31 kDa

Applications

Tested Applications:

Species Specificity:

human

Background Information

 $Proliferation of T \ lymphocytes \ is \ triggered \ by \ the \ interaction \ of \ IL-2 \ with \ its \ specific \ receptor \ following \ T$ lymphocyte activation. The receptor for IL-2 has three forms, generated by different combinations of three different proteins, the alpha chain (IL-2R alpha), the beta chain (IL-2R beta), and the gamma chain (IL-2R gamma) (PMID: 8476561). IL-2R alpha (also known as CD25) is a type I transmembrane protein present on activated T cells, activated B cells, some thymocytes, myeloid precursors, and oligodendrocytes. CD25 is also found on natural CD4+Foxp3+ Treg cells and a subset of human CD4+ memory T cells (PMID: 22585674).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1x10^6 PHA treated (3 days) human PBMCs were surface stained with CL750 Anti-Human CD3 and 5 ul Atlantic Blue™ Anti-Human CD25 (AB-65096, Clone:BC96) (red) or Atlantic Blue™ Mouse IgG1 Isotype Control (blue). 1x10^6 untreated human PBMCs were surface stained with 5 ul Atlantic Blue™ Anti-Human CD25 (AB-65096, Clone:BC96) (black dashed). Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Lymphocytes were gated.



1x10^6 PHA treated (3 days) human PBMCs were surface stained with CL750 Anti-Human CD3 and 5 ul Atlantic Blue™ Anti-Human CD25 (AB-65096, Clone: BC96) (right) or Atlantic Blue™ Mouse IgG1 Isotype Control (left). 1x10^6 untreated human PBMCs were surface stained with 5 ul Atlantic Blue™ Anti-Human CD25 (AB-65096, Clone:BC96) (center). Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Lymphocytes were gated.