

For Research Use Only

# APC Anti-Mouse CD8a (53-6.7)

Catalog Number: APC-65069

11 Publications



## Basic Information

Catalog Number:

APC-65069

Size:

100ug, 0.1 mg/ml

Source:

Rat

Isotype:

IgG2a, kappa

GenBank Accession Number:

BC030679

GeneID (NCBI):

12525

UNIPROT ID:

P01731

Full Name:

CD8 antigen, alpha chain

Purification Method:

Affinity purification

CloneNo.:

53-6.7

Excitation/Emission maxima wavelengths:

650 nm / 660 nm

## Applications

Tested Applications:

FC

Cited Applications:

IF, FC

Species Specificity:

mouse

Cited Species:

mouse

## Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8aβ heterodimer while CD8aa homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) γδ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).

## Notable Publications

Author	Pubmed ID	Journal	Application
Dingyi Wang	35236203	Drug Deliv	FC
Peixia Li	39447075	J Med Chem	
Yang Peng	39073014	Adv Healthc Mater	FC

## Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10<sup>6</sup> mouse splenocytes were surface co-stained with CoraLite® Plus 488 Anti-Mouse CD3 and 0.2 ug APC Anti-Mouse CD8a (APC-65069, Clone:53-6.7) or 0.2 ug Isotype Control. Cells were not fixed.