For Research Use Only

APC Anti-Mouse Foxp3 (3G3)

Catalog Number: APC-65089 3 Publications



Basic Information

Catalog Number:

APC-65089

Size:

100ug , 0.2 mg/ml

Source: Mouse

Isotype: IgG1, kappa GenBank Accession Number:

BC132333

GeneID (NCBI): 20371

UNIPROT ID: Q99JB6

Full Name: forkhead box P3 Purification Method:

Affinity purification

CloneNo.:

Excitation/Emission maxima

wavelengths: 650 nm / 660 nm

Applications

Tested Applications:

FC (Intra)

Cited Applications:

FC (Intra)

Species Specificity:

mouse

Cited Species:

mouse

Background Information

Foxp3 (forkhead box P3), also named as IPEX, JM2 and scurfin, is a member of the forkhead transcription factor family and is essential for normal immune homeostasis (PMID: 11138001; 20429413). Foxp3 is specifically expressed in CD4+ CD25+ regulatory T cells and is crucial for their development and suppressive activity (PMID: 12612578; 12612581). In scurfy mouse, disruption of Foxp3 results in the fatal lymphoproliferative disorder attributed to the lack of an immune-suppressive component in the immune system (PMID: 25683611; 11138001). Similar to mice, genetic defects in human Foxp3 are the cause of immunodeficiency polyendocrinopathy, enteropathy, X-linked syndrome (IPEX) which also known as X-linked autoimmunity-immunodeficiency syndrome (PMID: 11137993).

Notable Publications

Author	Pubmed ID	Journal	Application
Li Wu	35344716	Phytomedicine	FC (Intra)
Dingyi Wang	35236203	Drug Deliv	FC (Intra)
Yunkai Shang	37192052	ACS Nano	FC (Intra)

Storage


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1X10^6 BALB/c mouse splenocytes were surface stained with PE Anti-Mouse CD4 (GK1.5) and then fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set. Cells were then intracellularly stained with 0.06 ug APC Anti-Mouse Foxp3 (APC-65089, Clone:3G3) or 0.06 ug APC Mouse IgG1 Isotype Control (MOPC-21). It is recommended that the antibody dosage should not exceed 0.06ug.