

For Research Use Only

APC Anti-Mouse CD25/IL-2RA (PC61.5)

Catalog Number: APC-65137

3 Publications



Basic Information

Catalog Number:

APC-65137

Size:

100ug, 0.2 mg/ml

Source:

Rat

Isotype:

IgG1, lambda

GenBank Accession Number:

BC114437

GeneID (NCBI):

16184

UNIPROT ID:

P01590

Full Name:

interleukin 2 receptor, alpha chain

Purification Method:

Affinity purification

CloneNo.:

PC61.5

Excitation/Emission maxima wavelengths:

650 nm / 660 nm

Applications

Tested Applications:

FC

Cited Applications:

FC

Species Specificity:

mouse

Cited Species:

mouse

Background Information

Proliferation of T lymphocytes is triggered by the interaction of IL-2 with its specific receptor following T lymphocyte activation. The receptor for IL-2 has three forms, generated by different combinations of three different proteins, the alpha chain (IL-2R alpha), the beta chain (IL-2R beta), and the gamma chain (IL-2R gamma) (PMID: 8476561). IL-2R alpha (also known as CD25) is a type I transmembrane protein present on activated T cells, activated B cells, some thymocytes, myeloid precursors, and oligodendrocytes. CD25 is also found on natural CD4⁺Foxp3⁺ Treg cells (PMID: 22585674).

Notable Publications

Author	Pubmed ID	Journal	Application
Li Wu	35344716	Phytomedicine	FC
Tao Wang	38115100	J Neuroinflammation	FC
Zhuo Cheng	37544392	Acta Biomater	FC

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide and 0.5% BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



1x10⁶ untreated or anti-CD3/CD28 treated mouse splenocytes were surface stained with FITC Plus Anti-Mouse CD4 (GK1.5) (FITC-65104, Clone: GK1.5) and 0.1 ug APC Anti-Mouse CD25/IL-2RA (PC61.5) (APC-65137, Clone: PC61.5). Cells were not fixed.



1x10⁶ untreated or anti-CD3/CD28 treated (2 days) mouse splenocytes were co-surface stained with FITC Plus Anti-Mouse CD4 (GK1.5) and 0.25 ug APC Anti-Mouse CD25 (APC-65137, Clone:PC61.5) or 0.25 ug APC Rat IgG1 Isotype Control (HRPN). Cells were not fixed.



1x10⁶ anti-CD3/CD28 treated mouse splenocytes were surface stained with 0.1 ug APC Anti-Mouse CD25/IL-2RA (PC61.5) (APC-65137, Clone: PC61.5) (red) or APC Rat IgG1 Isotype Control (HRPN) (APC-65212, Clone: HRPN) (blue). 1x10⁶ untreated mouse splenocytes were surface stained with 0.1 ug APC Anti-Mouse CD25/IL-2RA (PC61.5) (APC-65137, Clone: PC61.5) (black, dashed). Cells were not fixed.