#### For Research Use Only

# APC Anti-Mouse CD279 (PD-1) (RMP1-30)



Catalog Number: APC-65142

1 Publications

**Basic Information** 

**Applications** 

Catalog Number:

APC-65142

100 $\mathrm{ug}$  , 0.2  $\mathrm{mg/ml}$ 

Rat Isotype:

IgG2b, kappa

**Tested Applications:** 

FC

**Cited Applications:** 

FC

Species Specificity:

Mouse

**Cited Species:** 

mouse

GenBank Accession Number:

BC119179 GeneID (NCBI):

Full Name:

18566

programmed cell death 1

**Purification Method:** 

Affinity purification

CloneNo.: RMP1-30

Excitation/Emission maxima

wavelengths: 650 nm / 660 nm

### **Background Information**

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Qing Zhang	35765095	Stem Cell Res Ther	FC


#### Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Phosphate based buffer with 0.09% sodium azide and 0.1% gelatin, pH 7.2.

## Selected Validation Data



1X10^6 unstimulated or anti-CD3/CD28 treated (2 days) BALB/C mouse splenocytes were surface stained with 0.2 ug APC Anti-Mouse CD279 (PD-1) (APC-65142, Clone: RMP1-30), and 0.5 ug FITC Plus Anti-Mouse CD3 (17A2) (FITC-65077, Clone: 17A2). Cells were not fixed.