For Research Use Only

APC Anti-Mouse PD-1/CD279 (RMP1-30)

Catalog Number: APC-65142 2 Publications




Basic Information	Catalog Number: APC-65142	GenBank Accession Number: BC119179		Purification Method: Affinity purification	
	Size: 100ug , 0.2 mg/ml	GenelD (NCI 18566	BI):	CloneNo.: RMP1-30	
	Source: Rat	Q02242 : Full Name:		Excitation/Emission maxima wavelengths: 650 nm / 660 nm	
	<mark>Isotype:</mark> IgG2b, kappa				
Applications	Tested Applications: FC				
	Cited Applications: FC				
	Species Specificity: mouse				
	Cited Species: mouse				
Background Information	Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).				
Notable Publications	Author	Pubmed ID	Journal	Арр	lication
	Qing Zhang	35765095	Stem Cell Res Ther	FC	
	Zhenzhen Xun	38956900	Adv Sci (Weinh)	FC	

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^6 untreated or anti-CD3/CD28 treated (2 days) mouse splenocytes were surface stained with 0.2 ug APC Anti-Mouse PD-1/CD279 (RMP1-30) (APC-65142, Clone: RMP1-30) and 0.2 ug APC Rat IgG2b Isotype Control (LTF-2) (APC-65211, Clone: LTF-2). Cells were not fixed.