For Research Use Only

APC Anti-Human CD73 (AD2)

Catalog Number: APC-65162



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Affinity purification

APC-65162

BC015940

GeneID (NCBI):

CloneNo.: AD2

wavelengths:

650 nm / 660 nm

100tests , 5 µl/test

4907 Full Name:

Excitation/Emission maxima

Mouse

5'-nucleotidase, ecto (CD73)

Calculated MW:

Isotype: IgG1, kappa

29 kDa, 63 kDa

Applications

Tested Applications:

Species Specificity:

Human

Background Information

CD73, also known as ecto-5'-nucleotidase (5'-NT), is a 70-kDa, glycosyl-phosphatidylinositol-linked membranebound glycoprotein found in most tissues (PMID: 18404475; 20179192). CD73 is an ectoenzyme that catalyzes the dephosphorylation of AMP and other nucleoside monophosphates (PMID: 9553767). In the human immune system, CD73 is expressed on subsets of T and B cells, on germinal center follicular dendritic cells, and on thymic medullary reticular fibroblasts and epithelial cells (PMID: 2137649; 9553767). CD73 is highly expressed in many human solid tumors and is closely involved in cancer progression (PMID: 20179192).

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1X10^6 human PBMCs were surface co-stained with CoraLite® Plus 405 CD3 and 5 ul APC Anti-Human CD73 (APC-65162, Clone:AD2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul APC Anti-Human CD73 (APC-65162, Clone:AD2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.