For Research Use Only

APC Anti-Rat CD8a (OX-8)

Catalog Number: APC-65180




Basic Information	Catalog Number: APC-65180	GenBank Accession Number: BC088126	Purification Method: Purified by protein-A affinity chromatography CloneNo.: OX-8 Excitation/Emission maxima wavelengths: 650 nm / 660 nm
	Size: 100ug , 200 µg/ml Source: Mouse Isotype: IgG1, kappa	GeneID (NCBI): 24930 UNIPROT ID: P07725 Full Name: CD8a molecule	
Background Information	CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8α or as a heterodimer of CD8α and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8αβ heterodimer while CD8αα homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) γδ T cells and NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T-cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).		
Storage	Storage: Store at 2-8°C. Avoid exposure a Storage Buffer: PBS with 0.1% sodium azide an	to light. Stable for one year after shipme d 0.5% BSA.	nt.

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 rat splenocytes were surface co-stained with PE Anti-Rat CD4 (OX-35) and 0.1 ug APC Anti-Rat CD8a (APC-65180, Clone: OX-8) or 0.1 ug APC Mouse IgG1 Isotype Control (MOPC-21). Cells were not fixed.