

For Research Use Only

# APC Anti-Human CD64 (10.1)

Catalog Number: APC-65253



## Basic Information

**Catalog Number:**

APC-65253

**Size:**

100 tests, 5 ul/test

**Source:**

Mouse

**Isotype:**

IgG1, kappa

**GenBank Accession Number:**

BC032634

**GeneID (NCBI):**

2209

**ENSEMBL Gene ID:**

ENSG00000150337

**UNIPROT ID:**

P12314

**Full Name:**

Fc fragment of IgG, high affinity Ia, receptor (CD64)

**Calculated MW:**

374 aa, 43 kDa

**Purification Method:**

N/A

**CloneNo.:**

10.1

**Excitation/Emission maxima wavelengths:**

650 nm / 660 nm

## Applications

**Tested Applications:**

FC

**Species Specificity:**

human

## Background Information

Fcγ receptor comprise a multigene family of integral membrane glycoproteins that exhibit complex activation or inhibitory effects on cell functions after aggregation by complexed immunoglobulin G (IgG) (PMID: 17005690). CD64, also known as Fcγ R1A, is a high-affinity receptor for the Fc region of IgG. It is expressed by monocytes/macrophages, activated neutrophils, dendritic cells, and early myeloid cells (PMID: 23293080; 19642859; 7680917). CD64 functions in both innate and adaptive immune responses.

## Storage

**Storage:**

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1x10<sup>6</sup> human PBMCs were surface stained with 5 ul APC Anti-Human CD64 (APC-65253, Clone:10.1) or APC Mouse IgG1 Isotype Control (APC-65124, Clone: MOPC-21). Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Monocytes were gated.



1x10<sup>6</sup> human PBMCs were surface stained with CL488 Anti-Human CD11b (CL488-65166) and 5 ul APC Anti-Human CD64 (APC-65253, Clone:10.1) or APC Mouse IgG1 Isotype Control (APC-65124, Clone: MOPC-21). Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Monocytes were gated.