

For Research Use Only

# FcZero-rAb™ APC Anti-Human IL-18R1/CD218a Rabbit Recombinant Antibody

Catalog Number: APC-FcA98150



## Basic Information

Catalog Number:

APC-FcA98150

Size:

100 tests, 5 ul/test

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC093975

GeneID (NCBI):

8809

Full Name:

interleukin 18 receptor 1

Calculated MW:

541 aa, 62 kDa

Purification Method:

Protein A purification

CloneNo.:

241673D3

Recommended Dilutions:

FC: 5 ul per 10<sup>6</sup> cells in a 100 µl suspension

Excitation/Emission maxima wavelengths:

650 nm / 660 nm

## Applications

Tested Applications:

FC

Species Specificity:

human

Positive Controls:

FC : human peripheral blood leukocytes,

## Background Information

IL-18R1 (CD218a) is a component of the interleukin-18 receptor (IL-18R) complex, which is critical for the signaling of IL-18, a pro-inflammatory cytokine. This receptor complex, consisting of IL-18R1 and the IL-18 receptor accessory protein (IL-18RAP), belongs to the immunoglobulin-like family of interleukin-1 receptors (PMID: 35461924). IL-18R1 plays a significant role in inducing the production of interferon-gamma (IFN-γ) and other inflammatory cytokines, contributing to innate and adaptive immune responses (PMID: 2550053). It is also suggested to play a role in hypertension and chronic kidney disease, where the IL-18/IL-18R1 signaling axis may have diagnostic and therapeutic potential (PMID: 35461924).

## Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1x10<sup>6</sup> human peripheral blood leukocytes were surface stained with FcZero-rAb™ PE Anti-Human NCAM1/CD56, and 5 ul APC Anti-Human IL-18R1/CD218a Rabbit RecAb (APC-FcA98150, Clone: 241673D3) or APC Rabbit IgG Isotype Control RecAb (APC-FcA98136, Clone: 240953C9). Cells were not fixed.