For Research Use Only

FcZero-rAb™ APC Anti-Mouse PDGFR beta/CD140b Rabbit Recombinant Antibody



Catalog Number: APC-FcA98207

Basic Information

Catalog Number:

APC-FcA98207

100ug , 200 $\mu g/ml$

Source:

Rabbit Isotype:

Immunogen Catalog Number:

GenBank Accession Number:

NM_008809.2

GeneID (NCBI): 18596

Full Name:

platelet derived growth factor

receptor, beta polypeptide

Calculated MW:

123kDa

Purification Method:

Protein A purification

CloneNo.: 241754F6

Recommended Dilutions:

FC: 0.1 ug per 10^6 cells in a 100 μ l

suspension

Excitation/Emission maxima

wavelengths: 650 nm / 660 nm

Applications

Tested Applications:

Species Specificity:

mouse

Positive Controls:

FC: NIH/3T3 cells,

Background Information

The Platelet-Derived Growth Factor Receptor Beta (PDGFR beta, also known as CD140b) is a receptor tyrosine kinase that plays a crucial role in various cellular processes, including cell proliferation, migration, and survival. In mice, PDGFR beta is particularly significant due to its expression in pericytes, which are cells that reside on the walls of capillaries and play a vital role in maintaining vascular stability and integrity (PMID: 20738866). In the central nervous system (CNS) of mice, PDGFR beta is predominantly expressed in pericytes and vascular smooth muscle cells. It is involved in the formation and maintenance of the blood-brain barrier, as well as in the regulation of angiogenesis and vascular stability. Research in mouse models has shown that PDGFR beta signaling is crucial for tissue repair and functional recovery after stroke. Mice with disrupted PDGFR beta signaling exhibit delayed recovery and increased infarction volume following cerebral ischemia (PMID: 21952111).

Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Selected Validation Data





.

1x10^6 NIH/3T3 cells (left) or mouse splenocytes (right) were surface stained with 0.1 ug APC Anti-Mouse PDGFR beta/CD140b Rabbit RecAb (APC-FcA98207, Clone: 241754F6) (red) or 0.1 ug APC Rabbit IgG Isotype Control RecAb (APC-FcA98136, Clone: 240953C9) (blue). Cells were not fixed.