

For Research Use Only

FcZero-rAb™ APC Anti-6*His, His-Tag Rabbit Recombinant Antibody

Catalog Number: APC-FcA98450



Basic Information

Catalog Number:

APC-FcA98450

Size:

100ug, 200 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

GeneID (NCBI):

UNIPROT ID:

HISTAG

Full Name:

Calculated MW:

1 kDa

Purification Method:

Protein A purification

CloneNo.:

251107A1

Recommended Dilutions:

FC (Intra): 0.1 ug per 10⁶ cells in a
100 µl suspension

Excitation/Emission maxima
wavelengths:

650 nm / 660 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

n/a

Positive Controls:

FC (Intra): Transfected HEK-293 cells,

Background Information

Protein tags are protein or peptide sequences located either on the C- or N- terminal of the target protein, which facilitates one or several of the following characteristics: solubility, detection, purification, localization and expression. His-tag is often used for affinity purification and binding assays. Expressed His-tagged proteins can be purified and detected easily because the string of histidine residues binds to several types of immobilized metal ions, including nickel, cobalt and copper, under specific buffer conditions. The His-tag antibody is a useful tool for monitoring of the His-tagged proteins, and recognizes His-tags placed at N-terminal, C-terminal, and internal regions of fusion proteins expressed in bacteria, insect, and mammalian cells.

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ HEK-293T or transfected HEK-293T cells were intracellularly stained with 0.1 ug APC Anti-6^{His}, His-Tag Rabbit RecAb (APC-FcA98450, Clone: 251107A1). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.

1x10⁶ transfected HEK-293T cells were intracellularly stained with 0.1 ug APC Anti-6^{His}, His-Tag Rabbit RecAb (APC-FcA98450, Clone: 251107A1) (red) or 0.1 ug APC Rabbit IgG Isotype Control RecAb (APC-FcA98136, Clone: 240953C9) (blue). 1x10⁶ untransfected HEK-293T cells were intracellularly stained with 0.1 ug APC Anti-6^{His}, His-Tag Rabbit RecAb (APC-FcA98450, Clone: 251107A1) (black, dashed). Cells were fixed with 4% PFA and permeabilized

