

For Research Use Only

# FcZero-rAb® APC-Cyanine7 Anti-Human CD64 (10.1) Rabbit Recombinant Antibody

Catalog Number:AY7-FcA65572



## Basic Information

**Catalog Number:**

AY7-FcA65572

**Size:**

100tests , 5 ul/test

**Source:**

Rabbit

**Isotype:**

IgG

**GenBank Accession Number:**

BC032634

**GeneID (NCBI):**

2209

**ENSEMBL Gene ID:**

ENSG00000150337

**UNIPROT ID:**

P12314

**Full Name:**

Fc fragment of IgG, high affinity Ia, receptor (CD64)

**Calculated MW:**

374 aa, 43 kDa

**Purification Method:**

Protein A purification

**CloneNo.:**

10.1

**Recommended Dilutions:**

FC: 5 ul per 10<sup>6</sup> cells in a 100 µl suspension

**Excitation/Emission maxima wavelengths:**

650 nm / 778 nm

## Applications

**Tested Applications:**

FC

**Species Specificity:**

human, monkey, non-human primates

**Positive Controls:**

FC : human PBMCs,

## Background Information

Fcγ receptor comprise a multigene family of integral membrane glycoproteins that exhibit complex activation or inhibitory effects on cell functions after aggregation by complexed immunoglobulin G (IgG) (PMID: 17005690). CD64, also known as Fcγ R1A, is a high-affinity receptor for the Fc region of IgG. It is expressed by monocytes/macrophages, activated neutrophils, dendritic cells, and early myeloid cells (PMID: 23293080; 19642859; 7680917). CD64 functions in both innate and adaptive immune responses.

## Storage

**Storage:**

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



1x10<sup>6</sup> human PBMCs were surface stained with CoraLite® Plus 488 Anti-Human CD11b (ICRF44), and 5 ul APC-Cyanine7 Anti-Human CD64 (10.1) Rabbit IgG RecAb (AY7-FcA65572, Clone: 10.1) or Rabbit IgG Isotype Control RecAb (AY7-FcA98136, Clone: 240953C9). Cells were incubated with MonoZero™ Monocytes blocking Reagent (PF00020) prior to staining. Cells were not fixed.



1x10<sup>6</sup> human PBMCs were surface stained with 5 ul APC-Cyanine7 Anti-Human CD64 (10.1) Rabbit IgG RecAb (AY7-FcA65572, Clone: 10.1) or Rabbit IgG Isotype Control RecAb (AY7-FcA98136, Clone: 240953C9). Cells were incubated with MonoZero™ Monocytes blocking Reagent (PF00020) prior to staining. Cells were not fixed.