FOR IN VITRO RESEARCH USE ONLY. NOT FOR USE IN HUMANS OR ANIMALS.

Recombinant human FARP1 protein



Basic Information Catalog Number: Ag21232 Form: Available lyophilized Species: human Expression Source: e coli-derived, PET28a, with N-terminal 6°His. Biological Activity: Not tested Endotoxin Level: Please contact the lab for more information Reconstitution: Reconstitution: Reconstitution and Storage Reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution.	VLDYVKEGGHKKVQFERKHSKIHSIRSLASQPTELNSE VLEQSQQSTSLTFGEGAESPGGQSCRRGKEPKVSAG EPGSHPSPAPRRSPAGNKQADGAASAPTEEEEEVVK DRTQQSKPQPPQPSTGSLTGSPHLSELSVNSQGGVA PANVTLSPNLSPDTKQASPLISPLLNDQACPRTDDEDE GRRKRFPTD (347-539 aa encoded by BC071592)
Available lyophilized Species: human Expression Source: e coli-derived, PET28a, with N-terminal 6*His. Biological Activity: Not tested Endotoxin Level: Please contact the lab for more information Reconstitution: Reconstitution with sterile water for short-term storage. After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	EPGSHPSPAPRRSPAGNKQADGAASAPTEEEEEVVK DRTQQSKPQPPQPSTGSLTGSPHLSELSVNSQGGVA PANVTLSPNLSPDTKQASPLISPLLNDQACPRTDDED GRRKRFPTD (347-539 aa encoded by BC071592) Shipping:
Species: human Expression Source: e coli-derived, PET28a, with N-terminal 6°His. Biological Activity: Not tested Endotoxin Level: Please contact the lab for more information Reconstitution: Reconstitute at 0.25 µg/µl in 200 µl sterile water for short-term storage. After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	DRTQQSKPQPPQPSTGSLTGSPHLSELSVNSQGGVA PANVTLSPNLSPDTKQASPLISPLLNDQACPRTDDED GRRKRFPTD (347-539 aa encoded by BC071592) Shipping:
Auman Expression Source: e coliderived, PET28a, with N-terminal 6*His. Biological Activity: Not tested Endotoxin Level: Please contact the lab for more information Reconstitution: Reconstitution: Reconstitution: Reconstitution with sterile water for short-term storage. After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	PANVTLSPNLSPDTKQASPLISPLLNDQACPRTDDEDE GRRKRFPTD (347-539 aa encoded by BC071592) Shipping:
Expression Source: e coli-derived, PET28a, with N-terminal 6*His.Biological Activity: Not testedEndotoxin Level: Please contact the lab for more informationReconstitution: Reconstitute at 0.25 µg/µl in 200 µl sterile water for short-term storage.After reconstitute at 0.25 µg/µl in 200 µl sterile water for short-term storage.After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details).If a different concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	GRRKRFPTD (347-539 aa encoded by BC071592) Shipping:
e coli-derived, PET28a, with N-terminal 6*His. Biological Activity: Not tested Endotoxin Level: Please contact the lab for more information Reconstitution: Reconstitute at 0.25 µg/µl in 200 µl sterile water for short-term storage. After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	(347-539 aa encoded by BC071592) Shipping:
Biological Activity: Not tested Endotoxin Level: Please contact the lab for more information Reconstitution: Reconstitute at 0.25 µg/µl in 200 µl sterile water for short-term storage. After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	Shipping:
Not tested Endotoxin Level: Please contact the lab for more information Reconstitution: Reconstitute at 0.25 µg/µl in 200 µl sterile water for short-term storage. After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	
Not tested Endotoxin Level: Please contact the lab for more information Reconstitution: Reconstitute at 0.25 µg/µl in 200 µl sterile water for short-term storage. After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	
Reconstitution and Storage Reconstitution: Reconstitute at 0.25 µg/µl in 200 µl sterile water for short-term storage. After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	
Reconstitution and StorageReconstitution: Reconstitute at 0.25 µg/µl in 200 µl sterile water for short-term storage. After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	
and Storage Reconstitute at 0.25 µg/µl in 200 µl sterile water for short-term storage. After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	
and Storage Reconstitute at 0.25 µg/µl in 200 µl sterile water for short-term storage. After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	The product is shipped at ambient temperature
After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	The product is shipped at ambient temperature.
on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	Upon receipt, store it immediately at the
equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	recommended temperature (see below).
If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	
adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	
ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	
the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	
gently pipet and wash down the sides of the vial to ensure full	
0,111	
90%, by SDS-PAGE with Coomassie Brilliant Blue staining.	
Formulation The purified protein was Lyophilized from sterile PBS (58mM N trehalose and 5 % mannitol are added as protectant before lyo imidazole.	• •
Store for up to 12 months at -20°C to -80°C as lyophilized power	ler.
Stability and Storage	
Storage of Short Term Storage:	
Aliquot and store at -20°C to -80°C for up to 3 months, reconstitution with sterile water and addition of an equal	
volume of glycerol. Avoid repeat freeze-thaw cycles.	
Selected Validation Data	
28 (₿a→	
43 kDa→	
28 kDa→ ←27 kDa	
20 kDa→	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522)E : proteintech@ptglab.com W:(toll free in USA), or 1(312) 455-8498ptglab.com(outside USA)(1312) 455-8498

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.